

Poulvac I L T

Introduction



Company name: [Fort Dodge Animal Health](#)
Address: **Flanders Road**
Hedge End
Southampton SO30 4QH
Telephone: **01489 781711**
Fax: **01489 788306**

Presentation

A freeze dried suspension of avian laryngotracheitis virus, Salisbury strain 146, 102.5 to 104.3EID50 per dose.

Uses

For the prevention of mortality and clinical signs due to infection with avian infectious laryngotracheitis virus. Onset of immunity is 14 days post vaccination. Duration of immunity has not been assessed, however, experience from field use indicates that once vaccinated birds remain immune to avian infectious laryngotracheitis virus infection during entire laying period.

Dosage and administration

Vaccine is administered to chickens from 4 weeks of age by eye drop.

1000 doses of freeze dried vaccine must be completely dissolved in 30ml of the sterile diluent provided and a drop of dispenser calibrated to deliver at least three hundredth of a millilitre per drop must be used. Remove the rubber stopper from the vaccine vial and add sterile diluent to half fill the vial. Replace the rubber stopper and shake gently so that all the vaccine material is completely dissolved.

Pour the dissolved vaccine into the remaining sterile diluent and shake again until completely mixed. The sterile diluent should be at room temperature when used. Fit the drop dispenser on the bottle. Hold the bird so that one eye is pointing upwards and allow one drop of vaccine to fall into the eye.

Correct administration:

Use clean vaccination materials.

Contact with the disinfectant makes the vaccine ineffective.

Protect the reconstituted vaccine from heat or direct sunlight.

Contra-indications, warnings, etc

Do not vaccinate diseased birds (except in case of an emergency vaccination).

Do not use in birds in lay.

No information is available on the safety and efficacy from the use of the product with any other. It is therefore recommended that no other vaccines than these should be administered within 14 days before or after vaccination with the product.

Do not mix with any other vaccine/immunological product.

In order to prevent spread of vaccine strain from vaccinated flocks to non-vaccinated flocks, all non-vaccinated birds present on the farm must be vaccinated at the same time.

Adverse reactions:

About four days after vaccination, redness and some swelling of the conjunctiva, or laboured breathing may be seen. Under good hygienic conditions, this reaction will not last longer than three days.

Withdrawal Period: Zero days.

Pharmaceutical precautions

Store and transport at +2°C to +8°C.

Protect from light. Do not freeze.

Use within two hours after reconstitution.

Keep out of reach and sight of children.

Legal category

POM-VPS (previously PML)

Packaging Quantities

Vaccine: packs of 10 vials containing 1,000 doses.

Diluent: packs of 10 x 30ml vials.

Further information

For Animal Treatment Only

Disposal:

Dispose of waste materials by boiling, incineration or immersion in an appropriate disinfectant approved for use by the competent authorities.

Marketing authorisation number

Vm 01596/4338